

Mathematical Physics

Individual Overall Contest

June 10, 2023

You can choose one of the following two questions.

Question 1. Consider a d -dimensional Ising model on a square lattice, which is periodic in all directions. The Hamiltonian is given by

$$H = - \sum_{a,b} J_{ab} \sigma_a \sigma_b - \sum_a B_a \sigma_a$$

where B_a is an external magnetic field, $\sigma_a = \pm 1$, and the spin-spin coupling $J_{ab} \equiv J(|\vec{a} - \vec{b}|)$ fall off with distance.

- (1) Compute the classical partition function when $J_{ab} = 0$ first, then show that for non-vanishing J , this can be written as

$$Z = \mathcal{N} \int \mathcal{D}\psi e^{-S[\psi]}, \quad S[\psi] = \sum_{a,b} \psi_a K_{ab} \psi_b - \sum_a \beta \psi_a B_a - \ln \cosh(2 \sum_b K_{ab} \psi_b)$$

where $K = (\beta J)^{-1}$

- (2) Show that at low temperature

$$S = \sum_{\mathbf{k}} [\phi_{\mathbf{k}} (c_0 + c_1 \mathbf{k} \cdot \mathbf{k}) \psi_{-\mathbf{k}} + c_2 \psi_{\mathbf{k}} B_{-\mathbf{k}}] + c_3 \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} \psi_{\mathbf{k}_1} \psi_{\mathbf{k}_2} \psi_{\mathbf{k}_3} \psi_{\mathbf{k}_4} \delta_{\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3 + \mathbf{k}_4} + \dots$$

where

$$\psi_a = \frac{1}{\sqrt{N}} \sum_{\mathbf{k}} e^{-i\mathbf{k} \cdot \mathbf{a}} \psi_{\mathbf{k}} \quad K_{ab} = \frac{1}{N} \sum_{\mathbf{k}} e^{-i\mathbf{k} \cdot (\mathbf{a} - \mathbf{b})} K_{\mathbf{k}}$$

then find coefficients c_i and explain when you can neglect the ellipsis in above expression.

- (3) For vanishing external magnetic field, upon Fourier transforming and rescaling, the action become a ϕ^4 theory

$$Z = Z_0 \int \mathcal{D}\phi e^{-S[\phi]} \quad S[\phi] = \int d^d x \left(\frac{1}{2} (\partial\phi)^2 + \frac{m}{2} \phi^2 + \lambda \phi^4 \right)$$

- (4) At what temperature does the Ising model have a transition? (HINT: the Ising model has a transition when m changes sign)

Question 2. Consider a free massless Dirac fermion in two-dimensions with Lagrangian

$$\mathcal{L} = i\bar{\psi}\not{\partial}\psi$$

The vector current $J_\mu \equiv \bar{\psi}\gamma_\mu\psi$ is conserved $\partial^\mu J_\mu = 0$. By the Poincaré lemma this means that we can introduce a scalar operator (field) $\phi(x)$ such that

$$J^\mu(x) = \epsilon^{\mu\nu}\partial_\nu\phi(x).$$

- (1) Answer to **either one** of the following two questions (they are equivalent, just stated in two different languages so that you can use the one which you feel more convenient):
 - (a) using the symmetries of the theory, write down the dynamical equations of the 2d Dirac theory as a field equation for the scalar field $\phi(x)$;
 - (b) compute the two-point function of $\phi(x)$ from the identity

$$\langle J^{\mu_1}(x_1) J^{\mu_2}(x_2) \rangle = \epsilon^{\mu_1\nu_1}\epsilon^{\mu_2\nu_2}\partial_{x_1^{\nu_1}}\partial_{x_2^{\nu_2}}\langle\phi(x_1)\phi(x_2)\rangle$$

[HINT: working in momentum space is more convenient]

- (2) can you write a local action for the field $\phi(x)$ whose variation yields the field equations obtained in (1)?
- (3) discuss briefly what your answer to (1) teaches us regarding the relation of bosons to fermions in two-dimensions
- (4) is your result (and discussion) consistent with the *Spin & Statistics* theorem of QFT?
- (5) Now couple your massless 2d Dirac fermion to an Abelian gauge field A_μ

$$\mathcal{L} = -\frac{1}{4e^2}F_{\mu\nu}F^{\mu\nu} + i\bar{\psi}\gamma^\mu(\partial_\mu - iA_\mu)\psi.$$

What is the particle spectrum of the resulting theory (\equiv massless 2d QED)? **To avoid all computation, the numerical value of the masses of the various particles is NOT required. Just say for each particle whether its mass is zero or non-zero.**